

PRIMER NOTE

Microsatellite loci for the field cricket, *Gryllus bimaculatus* and their cross-utility in other species of Orthoptera

DEBORAH A. DAWSON,* AMANDA J. BRETMAN*† and TOM TREGENZA†

*Department of Animal and Plant Sciences, University of Sheffield, Sheffield, S10 2TN, UK, †School of Biology, University of Leeds, Leeds, LS2 9JT, UK

Abstract

We have isolated 16 microsatellite loci in the field cricket, *Gryllus bimaculatus*. Nine loci were found to be polymorphic in *G. bimaculatus* and the number of alleles varied from seven to 14. All 16 loci were tested for amplification in nine other species. In the five species tested belonging to the same subfamily (Gryllinae), a minimum of nine loci amplified. These loci will be used to determine paternity as part of a study to investigate the genetic benefits of polyandry.

Keywords: cross-species amplification, field cricket, Gryllidae, *Gryllus bimaculatus*, microsatellite, Orthoptera

Received 3 September 2002; revision accepted 18 December 2002

The field or two-spot cricket, *Gryllus bimaculatus*, is an example of a species in which polyandry increases female reproductive success without obvious nongenetic benefits (Tregenza & Wedell 1998, 2002). In order to further investigate the genetic benefits of polyandry, we have isolated and characterized nine polymorphic microsatellite loci in this species.

DNA was extracted using a phenol : chloroform extraction method (Sambrook *et al.* 1989). For the production of the enriched microsatellite library, genomic DNA from a single male *G. bimaculatus* individual was digested with *Mbo*I (ABgene, Epsom, Surrey, UK) and enriched for (CA.GT)_n or (GA.CT)_n sequences (Amersham Pharmacia Biotech, Little Chalfont, Buckinghamshire, UK). The method used was essentially that described by Armour *et al.* (1994) with two modifications. To prevent duplicate clones, the DNA fragments were not PCR-amplified (polymerase chain reaction) before the enrichment hybridization. Secondly, a different plasmid cloning vector was used which was supplied pre-digested and dephosphorylated. The enriched fragments were ligated into pUC18-*Bam*HI/BAP (Amersham Pharmacia Biotech) and transformed into XL1-Blue competent cells (Stratagene,

La Jolly, CA, USA). Six hundred transformant clones were screened by hybridization to the sequences (CA.GT)_n or (GA.CT)_n radiolabelled with [α^{32} P]-dCTP, which identified 71 positives. Twenty-eight of the strongest positives were sequenced using BIG DYE Terminators on an ABI 377 Sequencer (Applied Biosystems, Foster City, CA, USA). Seventeen sequences were found to possess motifs with at least nine dinucleotide repeats and suitable for designing primers. These 17 sequences were confirmed unique using GENEJOCKEY Sequence Processor software (Biosoft, Cambridge, UK) and submitted to the EMBL database (Accession numbers: AJ315353–AJ315369). After submission to the EMBL database, the 17 sequences were double-checked unique using BLASTN 2.2.4 software (Altschul *et al.* 1997). All sequences were confirmed to be unique. Loci *Gbim10* and *Gbim12* (EMBL Accession numbers AJ315363 and AJ315365) are unique loci but exhibit 76% sequence homology between their respective flanking regions. The high number of base differences between the flanking sequence of *Gbim10* when compared with *Gbim12* suggests that these sequences are two different, unique loci and not different alleles of the same locus. The repeat region of *Gbim12* possesses additional (CG) motif repeats and fewer (CA) repeats. A different primer set was developed for each locus and different Orthoptera species were found to amplify with each primer set (Table 2), also suggesting these are two separate loci.

Correspondence: Deborah Dawson. Fax: + 44 (0)114 2220002; E-mail: d.a.dawson@sheffield.ac.uk

Table 1 Characterization of nine polymorphic microsatellites in the field cricket, *Gryllus bimaculatus*

Locus	EMBL no. and id	Repeat motif*	Primer sequence 5'-3' (and 5' primer label)	T_a (°C)	MgCl ₂ (mM)	Scored on	N	A	Expected allele size (bp)*	Observed allele size range (bp)	H_O	H_E
<i>Gbin04</i>	AJ315356	(GT) ₂₇	(F) (6-FAM)-CGACGATGTAGGCCCTGCGG (R) ATCCTACCAACACGGCAGCGG	65	1.5	ABI	15	11	241	205-239	0.80	0.89
<i>Gbin06</i>	AJ315359 CR2A05	(GT) ₂₅	(F) GCGATCGGAATCTTGAACCTGC (R) TTCCCTCGCCCTGACGACTCC	65	1.5	SS	15	7	204	160-220	0.80	0.83
<i>Gbin07</i>	AJ315360 CR2A12	(CG) ₃ (CA) ₁₀	(F) TCTTCCTTATCCCTATTCGCATCCC (R) CGGTGCCGACTTACAGTTG	65	1.0	SS	19	9	189	170-220	1.00	0.88
<i>Gbin08</i>	AJ315361 CR2B12	(CA) ₉ and (GA) ₃ AGAC(GA) ₂	(F) ACGTCAATACCATCAAGCCTTTCC (R) TCACCTTACAGGCCAACGCC	65	1.5	SS	20	9	176	170-210	0.85	0.86
<i>Gbin09</i>	AJ315362 CR2E08	(CA) ₉ CC(CA) ₄ CT(CA) ₂	(F) GATCTTCCCTTCTTCTTCTGTC (R) TCTCACCTCATATCCTGGG	60	1.0	SS	19	11	165	150-210	0.58	0.88
<i>Gbin11</i>	AJ315364 CR3A05	(CA) ₁₄	(F) (6-FAM)-CCCTTCCATATCCTACCCTCACC (R) CCAGATATCTTCTCCTGCGG	65	1.5	ABI	18	7	141	130-162	0.67	0.74
<i>Gbin13</i>	AJ315366 CR3E12	(GT) ₂₁ TT(GT) ₂	(F) TCCGGCACCAACTCTTAACTCAG (R) GATCAAACCCACCGGAACGTC	65	1.0	SS	16	9	246	200-310	0.88	0.86
<i>Gbin14</i>	AJ315367 CR3G03	(CA) ₇ TA(CA) ₁₁ CTA(CA) ₂ AG(CA) ₂	(F) (HEX)-GATCATTAATCCCTTCTTCTTGTTC (R) ATCCCTTCGCCGAGCTAAC	57	1.0	ABI	15	14	206	188-226	0.93	0.92
<i>Gbin15</i>	AJ315368 CR4E02	(CA) ₁₆	(F) GACTCGGGTACCCTTGTCC (R) ATCCGGAGCTTCAGCAAGGC	65	1.0	SS	20	8	179	150-220	0.95	0.87

PCR profile used: 94 °C 2 min, then 94 °C 1 min, T_a °C 30 s, 72 °C 30 s for 5 cycles, then 94 °C 30 s, T_a °C 30 s, 72 °C 30 s for 30 cycles, then 72 °C for 3 min.

*Of sequenced clone.

T_a , annealing temperature; SS, genotypes scored on 6% polyacrylamide gels stained with silver; ABI, genotypes scored on an ABI 377 Sequencer; N, number of individuals tested; A, number of alleles; H_O , observed heterozygosity; H_E , expected heterozygosity.

Seventeen primer pairs were designed with the assistance of PRIMER, version 3 (Rozen & Skaletsky 2000) and tested for amplification and polymorphism using 15–20 individuals. The *G. bimaculatus* individuals used to test for polymorphism were randomly selected from a laboratory population cultured in Leeds, which was originally established from individuals collected from the outskirts of Pretoria, South Africa. Each 10- μ L PCR contained 0.1–10 ng of genomic DNA, 1 μ M of each primer, 0.2 mM of each dNTP and 0.25 units *Taq* DNA polymerase (Thermoprime-Plus, ABgene) in the manufacturer's buffer [final concentrations 20 mM (NH₄)₂SO₄, 75 mM Tris-HCl pH 9.0, 0.01% (w/v) Tween], including 1.0 or 1.5 mM MgCl₂ (see Table 1). For two loci, PCR amplification was improved by the addition of 0.5% bovine serum albumin, fraction V (BSA; Sigma, St Louis, MO, USA) and 1% dimethyl sulphoxide (DMSO; Sigma), and by reducing the amount of template DNA to 1.0 ng or 0.1 ng. PCR amplification was performed in a Hybaid Touchdown thermal cycler (Thermo Hybaid, Ashford, Middlesex, UK). The PCR profile adopted is provided in Table 1. PCR amplification was confirmed on 2% agarose gels stained with ethidium bromide. Only one of the 17 primer sets, did not amplify a product (clone CR1F04, EMBL Accession number AJ315357).

Ten loci were characterized for polymorphism. Seven loci were analysed using 6% polyacrylamide gels stained with silver (Bassam *et al.* 1991; Promega, Southampton, UK) and three loci using an ABI 377 Sequencer. For the loci characterized on the ABI 377 Sequencer, the 5' end of the forward primer was labelled with a fluorescent phosphoramidite and gels analysed using GENESCAN software (version 3.1) and GENOTYPER DNA fragment analysis software (version 2.5). One locus was monomorphic (*Gbim12*, Table 2). Nine loci were polymorphic and these displayed between seven and 14 alleles in 15–20 *G. bimaculatus* individuals (Table 1). Primer sequences, expected PCR product sizes, allele size range and numbers of alleles for the nine polymorphic loci are given in Table 1. The observed and expected heterozygosity for each locus was calculated using CERVUS v2.0 (Marshall *et al.* 1998, 1999; Table 1). Only the locus (*Gbim09*) had a significant difference between observed and expected heterozygosity ($\chi^2 = 16.1$, $P < 0.001$). This could be due to the presence of a null allele.

Three loci, *Gbim04*, *Gbim11* and *Gbim14*, have been successfully used to assign paternity using an ABI 377 Sequencer (Bretman *et al.*, unpublished data).

The utility of all 16 *G. bimaculatus* loci in other species was assessed using three high quality DNA samples from each of nine other Orthopteran species (Table 2). The loci tested included: the nine characterized polymorphic loci, the monomorphic locus and six loci which amplified well and were polymorphic in *G. bimaculatus* but which were not characterized fully. EMBL Accession numbers and primer sequences for these loci are provided in Table 2.

The same PCR constituents were used as for *G. bimaculatus* but a different MgCl₂ concentration and PCR profile was adopted (Table 2).

Loci *Gbim04* and *Gbim06* jointly amplified in the highest number of other Orthopteran species (eight species). When tested in five species from the same subfamily, Gryllinae [genera: *Gryllus* (two species), *Teleogryllus*, *Acheta* and *Gryllodes*] between 9 and 14 loci amplified (56–88%). This high level of cross-amplification suggests these loci will be useful for genotyping in most of the 100–800 species belonging to the subfamily Gryllinae (species numbers dependent on the taxonomy used, Otte & Naskrecki 1997 or Hewitt 1979). Other studies have shown cross-species amplification of Orthopteran loci. Hockham *et al.* (1999) found 33–100% amplification of six bushcricket loci across 11 species of the same subfamily and King *et al.* (1998) found 50–100% amplification of four tree weta loci across four species in the same genus.

Four loci amplified in all five species of the subfamily Gryllinae but no other species. Within *Oecanthus nigricornis*, belonging to the same family as *G. bimaculatus* (Gryllidae) but a different subfamily (Oecanthinae), fewer loci (five) amplified. In species from three non-Gryllidae families a similar number of loci amplified (two to five loci; Table 2). More species need testing but this suggests that the genetic distance among species from different Gryllidae subfamilies is similar to the distance between species from different Orthopteran families. This may be due to the traditional classification, based upon phenotypic characteristics, not accurately reflecting phylogenetic relationships.

Acknowledgements

We thank Terry Burke for helpful advice and improvements to the manuscript, Jags Pandhal and Matt Bradshaw for assistance with colony screening and Martin Brinkhof, Dave Gray, Darryl Gwynne, Scott Sakaluk, Leigh Simmons and Mike Siva-Jothy for supplying Orthopteran DNA or tissue samples. Leon Hockham and Rob Buckland also kindly suggested improvements to the manuscript. This work was performed at the Sheffield Molecular Genetics Facility supported by the Natural Environment Research Council.

References

- Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research*, **25**, 3389–3402.
- Armour JAL, Neumann R, Gobert S, Jeffreys AJ (1994) Isolation of human simple repeat loci by hybridization selection. *Human Molecular Genetics*, **3**, 599–605.
- Bassam BJ, Caetano-Anolles G, Gresshoff PM (1991) Fast and sensitive silver staining of polyacrylamide gels. *Analytical Biochemistry*, **196**, 80.

Table 2 Cross-amplification of 16 *Gryllus bimaculatus* microsatellite loci in nine other species belonging to the order Orthoptera

Locus	EMBL no. and id.	Repeat motif*	Primer sequence (5'-3')	PCR profile (mm)	MgCl ₂ (mm)	Species†	Family											Total other Orthoptera spp. (/9)	
							Subfamily					No. of spp. amplifying per locus							
							Gr	Gr	Gr	Gr	Gr	na	H	na	na	A	T		P
Gbim01	AJ15353	(CA) ₂₄	(F) ATTTGGCCAAAGCATAGCAGC (R) AATGAGATATCCAGCGGGC	65.td	2.0	+	+	+	+	+	+	+	-	+	+	-	5	2	7
Gbim02	AJ15354	(CA) ₁₀	(F) GATCCTTCCCTCCATTTTCCATC (R) CCGACTGGTCTCCCGACT	60.td	0.6	+	+	+	+	+	+	+	-	+	-	-	5	1	6
Gbim03	AJ15355	(CT) ₃₁ (CA) ₁₇	(F) GCGAATCCAGAGAGTACCC (R) AGACAGACCCGCTACACCCG	65.td	1.0	+	+	+	-	+	-	-	-	+	-	-	3	0	3
Gbim04	AJ15356	(GT) ₂₇	(F) CGACGTATGAGCCCTGGG (R) ATCTTACCAAACACGGCAGG	65.td	1.5	+,P	+	+	+	+	+	+	+	+	+	+	5	3	8
Gbim05	AJ15358	(GT) ₁₂ (GTA) ₇	(F) CCGGATGTGGTAGAGTAG (R) TATCAAACGGCCACGGAAC	65.td	0.6	+	+	+	+	+	+	+	-	-	-	-	5	0	5
Gbim06	AJ15359	(GT) ₂₅	(F) GCGATGCGAATCTTGAATGC (R) TTCTCGCCTTGAGACTCC	65.td	1.5	+,P	+	+	+	+	+	+	+	+	+	+	6	2	8
Gbim07	AJ15360	(CG) ₃ (CA) ₁₀	(F) TCTTCTTATCCTTATTCGCATCCC (R) CGGTGCCGACTTACACGTTG	65.td	1.0	+,P	+	+	+	+	+	+	-	-	-	-	5	0	5
Gbim08	AJ15361	(CA) ₉ and (GA) ₃	(F) AGGTCAATACCATCAAGCCTTTCC (R) TCACCTTACAGGGCCAAAGCC	65.td	1.5	+,P	+	+	+	+	+	+	+	+	+	-	5	1	6
Gbim09	AJ15362	(CA) ₉ CC(CA) ₄ CT	(F) GATCTTCTTCTTCTTCTTCTGTC (R) TCTCACCTCATATCTCTGGG	60.td	0.6	+,P	-	-	-	-	-	-	-	-	-	-	0	0	0
Gbim10†	AJ15363	(CA) ₂₁	(F) ACACCAGGGGAAATGCTGGAG (R) CCATCCAGACAGAAACGAGC	60.td	1.0	+	+	+	+	+	+	+	-	-	-	-	5	0	5
Gbim11	AJ15364	(CA) ₁₄	(F) CCTTCCATATCTTACCTCACC (R) CCAGATATCTTCTCCTCGG	65.td	1.5	+,P	+	+	+	+	+	+	+	+	+	+	5	0	5
Gbim12†	AJ15365	(CG) ₄ (CA) ₁₂	(F) ATCATCACCAGCAATGCC (R) ACCATGTACGCTTGGGAG	65.td	0.6	+,M	-	-	-	-	-	-	-	-	-	-	0	0	0
Gbim13	AJ15366	(GT) ₂₁ TT(GT) ₂	(F) TCCGGACCAACTTTAAGTCAG (R) GATCAACACCAGGAAGTC	65.td	1.0	+,P	+	+	+	+	+	+	-	-	-	-	3	0	3
Gbim14	AJ15367	(CA) ₇ TA(CA) ₁₁ CTA	(F) GATCAATTATCTTCTTCTGTTTC (R) ATCCTTCGGCGAGCTAAC	60.td	1.0	+,P	+	+	+	+	+	+	-	-	-	-	4	0	4

Table 2 Continued

Locus	EMBL no. and id. motif*	Repeat motif*	Primer sequence (5'–3')	PCR profile (mm)	Family												
					Gr	Gr	Gr	Gr	O	na	H	na	A	T	P		
Gbim15	AJ315368 CR4E02	(CA) ₁₆	(F) GACTCGGGTACCCCTTCTCG (R) ATCCGGAGCTTCAGCAAGGC	65.td	+	+	+	-	+	+	-	+	-	5	1	6	
Gbim16	AJ315369 CR6B11	(CGG) ₄ and (GAGCG) ₅	(F) GACGGCCCGGAATCACTAG (R) CCAGCCCGGAAGCACGAG	65.td	+	+	+	-	+	-	-	+	-	4	1	5	
Total number of loci amplifying (/16)					16	14	14	13	10	9	5	5	4	4	2		

PCR profiles used:

65.td = 94 °C for 2 min for one cycle, then 94 °C for 30 s, X °C for 30 s, 72 °C for 30 s for two cycles (where X = 65 °C then 63, 61, 59, 57) then 94 °C for 30 s, 55 °C for 30 s, 72 °C for 30 s for 30 cycles, and finally 72 °C for 3 min.

60.td = 94 °C for 2 min for one cycle, then 94 °C for 30 s, 72 °C for 30 s, 72 °C for 30 s for two cycles (where Y = 60 °C then 58, 56, 54, 52) then 94 °C for 30 s, 50 °C for 30 s, 72 °C for 30 s for 30 cycles, and finally 72 °C for 3 min.

*Of sequenced clone.

†Three unrelated individuals of each species were tested and species are ordered as most closely related to *Gryllus bimaculatus* (based on Gwynne 1995; Huang *et al.* 2000).

‡The sequences of Gbim10 and Gbim12 (AJ315363 and AJ315365) exhibit 76% sequence homology in their respective flanking regions (see text).

na, not applicable.

G, Gryllidae family; A, Anostoptmatidae family; T, Tettigoniidae family; P, Prophalangopsidae family; Gr, Gryllinae Subfamily; O, Oecanthinae subfamily; H, Haglidae subfamily.

GB, *Gryllus bimaculatus*, field cricket (origin of loci); GC, *Gryllus campestris*, field cricket (*G. campestris* is the sister species of *G. bimaculatus*); GT, *Gryllus texensis*, field cricket; TO, *Teleogryllus oceanicus*, field cricket; AD, *Acheta domestica*, black house cricket; CS, *Gryllodes sigillatus*, tropical house cricket; ON, *Oecanthus nigricornis*, black horned tree cricket; HF, *Hemianthus furcifer*, ground weta; CN, *Conocephalus nigropleurum*, black sided meadow katydid; CS, *Cyphoderris strepitans*, sage bush cricket.

+, amplification; -, no amplification; P, polymorphic (see Table 1); M, monomorphic.

6 PRIMER NOTE

- Gwynne DT (1995) Phylogeny of the Ensifera (Orthoptera): a hypothesis supporting multiple origins of acoustical signalling, complex spermatophores and maternal care in crickets, katydids, and weta. *Journal of Orthoptera Research*, **4**, 203–218.
- Hewitt GM (1979) *Animal Cytogenetics 3, Insecta 1, Orthoptera: Grasshoppers and Crickets*. Gerbruder Borntrager, Berlin, Germany.
- Hockham LR, Graves JA, Ritchie MG (1999) Isolation and characterization of microsatellite loci in the bushcricket *Ephippiger ephippiger* (Orthoptera: Tettigoniidae). *Molecular Ecology*, **8**, 905–906.
- Huang Y, Orti G, Sutherlin M, Siegel-Causey D, Duhachek A, Zera AJ (2000) Phylogenetic relationships of North American field crickets inferred from mitochondrial DNA data. *Molecular Phylogenetics and Evolution*, **17**, 48–57.
- King TM, Hanotte O, Burke T, Wallis GP (1998) Characterisation of four microsatellite loci in tree weta (Orthoptera: Stenopelmataidae): their potential usefulness for the study of *Hemideina*. *Molecular Ecology*, **7**, 663–664.
- Marshall TC, Slate J, Kruuk LEB, Pemberton JM (1998, 1999) Statistical confidence for likelihood-based paternity inference in natural populations. *Molecular Ecology*, **7**, 639–655. <http://helios.bto.ed.ac.uk/evolgen/cervus/cervus.html>
- Otte D, Naskrecki P (1997) Orthoptera species file online. <http://140.247.119.145/Orthoptera/>.
- Rozen S, Skaletsky HJ (2000) PRIMER3 on the WWW for general users and for biologist programmers. In: *Bioinformatics Methods and Protocols: Methods in Molecular Biology* (eds Krawetz S, Misener S) pp. 365–386. Humana Press, Totowa, NJ. http://www-genome.wi.mit.edu/genome_software/other/primer3.html.
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular Cloning. A Laboratory Manual*, 2nd edn. Cold Spring Harbor Laboratory Press, New York, USA.
- Tregenza T, Wedell N (1998) Benefits of multiple mates in the cricket *Gryllus bimaculatus*. *Evolution*, **52**, 1726–1730.
- Tregenza T, Wedell N (2002) Polyandrous females avoid costs of inbreeding. *Nature*, **415**, 71–73.